

14), C180 (SEQ ID NO: 15), C204 (SEQ ID NO: 16), C209 (SEQ ID NO: 17), C266 (SEQ ID NO: 18), C342 (SEQ ID NO: 19), and C403 (SEQ ID NO: 20).

Please replace the paragraph beginning on page 6, line 5, and ending on page 8, line 22, with the following replacement paragraph:

The hypersensitive response elicitor polypeptide or protein from *Erwinia chrysanthemi* has an amino acid sequence corresponding to SEQ ID NO: 21 as follows:

Met Gln Ile Thr Ile Lys Ala His Ile Gly Gly Asp Leu Gly Val Ser  
1 5 10 15  
Gly Leu Gly Ala Gln Gly Leu Lys Gly Leu Asn Ser Ala Ala Ser Ser  
20 25 30  
Leu Gly Ser Ser Val Asp Lys Leu Ser Ser Thr Ile Asp Lys Leu Thr  
35 40 45  
Ser Ala Leu Thr Ser Met Met Phe Gly Gly Ala Leu Ala Gln Gly Leu  
50 55 60  
Gly Ala Ser Ser Lys Gly Leu Gly Met Ser Asn Gln Leu Gly Gln Ser  
65 70 75 80  
Phe Gly Asn Gly Ala Gln Gly Ala Ser Asn Leu Leu Ser Val Pro Lys  
85 90 95  
Ser Gly Gly Asp Ala Leu Ser Lys Met Phe Asp Lys Ala Leu Asp Asp  
100 105 110  
Leu Leu Gly His Asp Thr Val Thr Lys Leu Thr Asn Gln Ser Asn Gln  
115 120 125  
Leu Ala Asn Ser Met Leu Asn Ala Ser Gln Met Thr Gln Gly Asn Met  
130 135 140  
Asn Ala Phe Gly Ser Gly Val Asn Asn Ala Leu Ser Ser Ile Leu Gly  
145 150 155 160  
Asn Gly Leu Gly Gln Ser Met Ser Gly Phe Ser Gln Pro Ser Leu Gly  
165 170 175  
Ala Gly Gly Leu Gln Gly Leu Ser Gly Ala Gly Ala Phe Asn Gln Leu  
180 185 190  
Gly Asn Ala Ile Gly Met Gly Val Gly Gln Asn Ala Ala Leu Ser Ala  
195 200 205  
Leu Ser Asn Val Ser Thr His Val Asp Gly Asn Asn Arg His Phe Val  
210 215 220  
Asp Lys Glu Asp Arg Gly Met Ala Lys Glu Ile Gly Gln Phe Met Asp  
225 230 235 240

A2  
Cool'd

Gln	Tyr	Pro	Glu	Ile	Phe	Gly	Lys	Pro	Glu	Tyr	Gln	Lys	Asp	Gly	Trp	245	250	255	
Ser	Ser	Pro	Lys	Thr	Asp	Asp	Lys	Ser	Trp	Ala	Lys	Ala	Leu	Ser	Lys	260	265	270	
Pro	Asp	Asp	Asp	Gly	Met	Thr	Gly	Ala	Ser	Met	Asp	Lys	Phe	Arg	Gln	275	280	285	
Ala	Met	Gly	Met	Ile	Lys	Ser	Ala	Val	Ala	Gly	Asp	Thr	Gly	Asn	Thr	290	295	300	
Asn	Leu	Asn	Leu	Arg	Gly	Ala	Gly	Gly	Ala	Ser	Leu	Gly	Ile	Asp	Ala	305	310	315	320
Ala	Val	Val	Gly	Asp	Lys	Ile	Ala	Asn	Met	Ser	Leu	Gly	Lys	Leu	Ala	325	330	335	
Asn	Ala																		

This hypersensitive response elicitor polypeptide or protein has a molecular weight of 34 kDa, is heat stable, has a glycine content of greater than 16%, and contains substantially no cysteine. The *Erwinia chrysanthemi* hypersensitive response elicitor polypeptide or protein is encoded by a DNA molecule having a nucleotide sequence corresponding to SEQ ID NO: 22 as follows:

CGATTTTACC	CGGGTGAACG	TGCTATGACC	GACAGCATCA	CGGTATTCTGA	CACCGTTACG	60
GCGTTTATGG	CCGCGATGAA	CCGGCATCAG	GCGGCGCGCT	GGTCGCCGCA	ATCCGGCGTC	120
GATCTGGTAT	TTCAGTTTGG	GGACACCGGG	CGTGAACCTCA	TGATGCAGAT	TCAGCCGGGG	180
CAGCAATATC	CCGGCATGTT	GCGCACGCTG	CTCGCTCGTC	GTTATCAGCA	GGCGGCAGAG	240
TGCGATGGCT	GCCATCTGTG	CCTGAACGGC	AGCGATGTAT	TGATCCTCTG	GTGGCCGCTG	300
CCGTCGGATC	CCGGCAGTTA	TCCGCAGGTG	ATCGAACGTT	TGTTTGAACT	GGCGGGAATG	360
ACGTTGCCGT	CGCTATCCAT	AGCACCGACG	GCGCGTCCGC	AGACAGGGAA	CGGACGCGCC	420
CGATCATTA	GATAAAGGCG	GCTTTTTTTA	TTGCAAAACG	GTAACGGTGA	GGAACCGTTT	480
CACCGTCGGC	GTCACCTAGT	AACAAGTATC	CATCATGATG	CCTACATCGG	GATCGGCGTG	540
GGCATCCGTT	GCAGATACTT	TTGCGAACAC	CTGACATGAA	TGAGGAAACG	AAATTATGCA	600
AATTACGATC	AAAGCGCACA	TCGGCGGTGA	TTTGGGCGTC	TCCGGTCTGG	GGCTGGGTGC	660
TCAGGGACTG	AAAGGACTGA	ATTCCGCGGC	TTCATCGCTG	GGTTCAGCG	TGGATAAACT	720
GAGCAGCACC	ATCGATAAGT	TGACCTCCGC	GCTGACTTCG	ATGATGTTTG	GCGGCGCGCT	780
GGCGCAGGGG	CTGGGCGCCA	GCTCGAAGGG	GCTGGGGATG	AGCAATCAAC	TGGGCCAGTC	840
TTTCGGCAAT	GGCGCGCAGG	GTGCGAGCAA	CCTGCTATCC	GTACCGAAAT	CCGGCGGCGA	900

AZ  
Cont'd

TGCGTTGTCA AAAATGTTTG ATAAAGCGCT GGACGATCTG CTGGGTCATG ACACCGTGAC	960
CAAGCTGACT AACCAGAGCA ACCAACTGGC TAATTCAATG CTGAACGCCA GCCAGATGAC	1020
CCAGGGTAAT ATGAATGCGT TCGGCAGCGG TGTGAACAAC GCACTGTCGT CCATTCTCGG	1080
CAACGGTCTC GGCCAGTCGA TGAGTGGCTT CTCTCAGCCT TCTCTGGGGG CAGGCGGCTT	1140
GCAGGGCCTG AGCGGCGCGG GTGCATTCAA CCAGTTGGGT AATGCCATCG GCATGGGCGT	1200
GGGGCAGAAT GCTGCGCTGA GTGCGTTGAG TAACGTCAGC ACCCACGTAG ACGGTAACAA	1260
CCGCCACTTT GTAGATAAAG AAGATCGCGG CATGGCGAAA GAGATCGGCC AGTTTATGGA	1320
TCAGTATCCG GAAATATTCG GTAAACCGGA ATACCAGAAA GATGGCTGGA GTTCGCCGAA	1380
GACGGACGAC AAATCCTGGG CTAAAGCGCT GAGTAAACCG GATGATGACG GTATGACCGG	1440
CGCCAGCATG GACAAATTCC GTCAGGCGAT GGGTATGATC AAAAGCGCGG TGGCGGGTGA	1500
TACCGGCAAT ACCAACCTGA ACCTGCGTGG CGCGGGCGGT GCATCGCTGG GTATCGATGC	1560
GGCTGTCTGTC GGCATAAAA TAGCCAACAT GTCGCTGGGT AAGCTGGCCA ACGCCTGATA	1620
ATCTGTGCTG GCCTGATAAA GCGGAAACGA AAAAAGAGAC GGGGAAGCCT GTCTCTTTTC	1680
TTATTATGCG GTTTATGCGG TTACCTGGAC CGGTTAATCA TCGTCATCGA TCTGGTACAA	1740
ACGCACATTT TCCCGTTCAT TCGCGTCGTT ACGCGCCACA ATCGCGATGG CATCTTCCTC	1800
GTCGCTCAGA TTGCGCGGCT GATGGGGAAC GCCGGTGGA ATATAGAGAA ACTCGCCGGC	1860
CAGATGGAGA CACGTCTGCG ATAAATCTGT GCCGTAACGT GTTCTATCC GCCCCTTAG	1920
CAGATAGATT GCGGTTTCGT AATCAACATG GTAATGCGGT TCCGCCTGTG CGCCGGCCGG	1980
GATCACCACA ATATTCATAG AAAGCTGTCT TGCACCTACC GTATCGCGGG AGATACCGAC	2040
AAAATAGGGC AGTTTTTGCG TGGTATCCGT GGGGTGTTCC GGCCTGACAA TCTTGAGTTG	2100
GTTTCGTCATC ATCTTTCTCC ATCTGGGCGA CCTGATCGGT T	2141

Please replace the paragraph beginning on page 8, line 25, and ending on page 11, line 8, with the following replacement paragraph:

The hypersensitive response elicitor polypeptide or protein derived from *Erwinia amylovora* has an amino acid sequence corresponding to SEQ ID NO: 23 as follows:

A3

Met	Ser	Leu	Asn	Thr	Ser	Gly	Leu	Gly	Ala	Ser	Thr	Met	Gln	Ile	Ser
1				5					10					15	
Ile	Gly	Gly	Ala	Gly	Gly	Asn	Asn	Gly	Leu	Leu	Gly	Thr	Ser	Arg	Gln
			20					25						30	

Asn Ala Gly Leu Gly Gly Asn Ser Ala Leu Gly Leu Gly Gly Gly Asn  
 35 40 45

Gln Asn Asp Thr Val Asn Gln Leu Ala Gly Leu Leu Thr Gly Met Met  
 50 55 60

Met Met Met Ser Met Met Gly Gly Gly Gly Leu Met Gly Gly Gly Leu  
 65 70 75 80

Gly Gly Gly Leu Gly Asn Gly Leu Gly Gly Ser Gly Gly Leu Gly Glu  
 85 90 95

Gly Leu Ser Asn Ala Leu Asn Asp Met Leu Gly Gly Ser Leu Asn Thr  
 100 105 110

Leu Gly Ser Lys Gly Gly Asn Asn Thr Thr Ser Thr Thr Asn Ser Pro  
 115 120 125

Leu Asp Gln Ala Leu Gly Ile Asn Ser Thr Ser Gln Asn Asp Asp Ser  
 130 135 140

Thr Ser Gly Thr Asp Ser Thr Ser Asp Ser Ser Asp Pro Met Gln Gln  
 145 150 155 160

Leu Leu Lys Met Phe Ser Glu Ile Met Gln Ser Leu Phe Gly Asp Gly  
 165 170 175

Gln Asp Gly Thr Gln Gly Ser Ser Ser Gly Gly Lys Gln Pro Thr Glu  
 180 185 190

Gly Glu Gln Asn Ala Tyr Lys Lys Gly Val Thr Asp Ala Leu Ser Gly  
 195 200 205

Leu Met Gly Asn Gly Leu Ser Gln Leu Leu Gly Asn Gly Gly Leu Gly  
 210 215 220

Gly Gly Gln Gly Gly Asn Ala Gly Thr Gly Leu Asp Gly Ser Ser Leu  
 225 230 235 240

Gly Gly Lys Gly Leu Gln Asn Leu Ser Gly Pro Val Asp Tyr Gln Gln  
 245 250 255

Leu Gly Asn Ala Val Gly Thr Gly Ile Gly Met Lys Ala Gly Ile Gln  
 260 265 270

Ala Leu Asn Asp Ile Gly Thr His Arg His Ser Ser Thr Arg Ser Phe  
 275 280 285

Val Asn Lys Gly Asp Arg Ala Met Ala Lys Glu Ile Gly Gln Phe Met  
 290 295 300

Asp Gln Tyr Pro Glu Val Phe Gly Lys Pro Gln Tyr Gln Lys Gly Pro  
 305 310 315 320

Gly Gln Glu Val Lys Thr Asp Asp Lys Ser Trp Ala Lys Ala Leu Ser  
 325 330 335

Lys Pro Asp Asp Asp Gly Met Thr Pro Ala Ser Met Glu Gln Phe Asn  
 340 345 350  
 Lys Ala Lys Gly Met Ile Lys Arg Pro Met Ala Gly Asp Thr Gly Asn  
 355 360 365  
 Gly Asn Leu Gln Ala Arg Gly Ala Gly Gly Ser Ser Leu Gly Ile Asp  
 370 375 380  
 Ala Met Met Ala Gly Asp Ala Ile Asn Asn Met Ala Leu Gly Lys Leu  
 385 390 395 400  
 Gly Ala Ala

This hypersensitive response elicitor polypeptide or protein has a molecular weight of about 39 kDa, has a pI of approximately 4.3, and is heat stable at 100°C for at least 10 minutes. This hypersensitive response elicitor polypeptide or protein has substantially no cysteine. The hypersensitive response elicitor polypeptide or protein derived from *Erwinia amylovora* is more fully described in Wei, Z.-M., R. J. Laby, C. H. Zumoff, D. W. Bauer, S.-Y. He, A. Collmer, and S. V. Beer, "Harpin, Elicitor of the Hypersensitive Response Produced by the Plant Pathogen *Erwinia amylovora*," *Science* 257:85-88 (1992), which is hereby incorporated by reference. The DNA molecule encoding this polypeptide or protein has a nucleotide sequence corresponding to SEQ ID NO: 24 as follows:

```

AAGCTTCGGC ATGGCACGTT TGACCGTTGG GTCGGCAGGG TACGTTTGAA TTATTCATAA      60
GAGGAATACG TTATGAGTCT GAATACAAGT GGGCTGGGAG CGTCAACGAT GCAAATTTCT      120
ATCGGCGGTG CGGGCGGAAA TAACGGGTTG CTGGGTACCA GTCGCCAGAA TGCTGGGTTG      180
GGTGGCAATT CTGCACTGGG GCTGGGCGGC GGTAATCAAA ATGATACCGT CAATCAGCTG      240
GCTGGCTTAC TCACCGGCAT GATGATGATG ATGAGCATGA TGGGCGGTGG TGGGCTGATG      300
GGCGGTGGCT TAGGCGGTGG CTTAGGTAAT GGCTTGGGTG GCTCAGGTGG CCTGGGCGAA      360
GGACTGTCGA ACGCGCTGAA CGATATGTTA GGCGGTTTCG TGAACACGCT GGGCTCGAAA      420
GGCGGCAACA ATACCACTTC AACAAACAAT TCCCCGCTGG ACCAGGCGCT GGGTATTAAC      480
TCAACGTCCC AAAACGACGA TTCCACCTCC GGCACAGATT CCACCTCAGA CTCCAGCGAC      540
CCGATGCAGC AGCTGCTGAA GATGTTTCAGC GAGATAATGC AAAGCCTGTT TGGTGATGGG      600
CAAGATGGCA CCCAGGGCAG TTCTCTTGGG GGCAAGCAGC CGACCGAAGG CGAGCAGAAC      660
GCCTATAAAA AAGGAGTCAC TGATGCGCTG TCGGGCCTGA TGGGTAATGG TCTGAGCCAG      720
CTCCTTGGCA ACGGGGGACT GGGAGGTGGT CAGGGCGGTA ATGCTGGCAC GGGTCTTGAC      780
GGTTCGTCGC TGGGCGGCAA AGGGCTGCAA AACCTGAGCG GGCCGGTGGA CTACCAGCAG      840
TTAGGTAACG CCGTGGGTAC CGGTATCGGT ATGAAAGCGG GCATTCAGGC GCTGAATGAT      900
  
```

ATCGGTACGC ACAGGCACAG TTCAACCCGT TCTTTCGTCA ATAAAGGCGA TCGGGCGATG 960  
 GCGAAGGAAA TCGGTCAGTT CATGGACCAG TATCCTGAGG TGTTTGCAA GCCGCAGTAC 1020  
 CAGAAAGGCC CGGGTCAGGA GGTGAAAACC GATGACAAAT CATGGGCAA AGCACTGAGC 1080  
 AAGCCAGATG ACGACGGAAT GACACCAGCC AGTATGGAGC AGTTCAACAA AGCCAAGGGC 1140  
 ATGATCAAAA GGCCCATGGC GGGTGATACC GGCAACGGCA ACCTGCAGGC ACGCGGTGCC 1200  
 GGTGGTTCTT CGCTGGGTAT TGATGCCATG ATGGCCGGTG ATGCCATTAA CAATATGGCA 1260  
 CTTGGCAAGC TGGGCGCGGC TTAAGCTT 1288

Please replace the paragraph beginning on page 11, line 11, and ending on page 13, line 47, with the following replacement paragraph:

Another potentially suitable hypersensitive response elicitor from *Erwinia amylovora* is disclosed in U.S. Patent Application Serial No. 09/120,927, which is hereby incorporated by reference. The protein is encoded by a DNA molecule having a nucleic acid sequence of SEQ ID NO: 25 as follows:

ATGTCAATTC TTACGCTTAA CAACAATACC TCGTCCTCGC CGGGTCTGTT CCAGTCCGGG 60  
 GGGGACAACG GGCTTGGTGG TCATAATGCA AATTCTGCGT TGGGGCAACA ACCCATCGAT 120  
 CGGCAAACCA TTGAGCAAAT GGCTCAATTA TTGGCGGAAC TGTTAAAGTC ACTGCTATCG 180  
 CCACAATCAG GTAATGCGGC AACCGGAGCC GGTGGCAATG ACCAGACTAC AGGAGTTGGT 240  
 AACGCTGGCG GCCTGAACGG ACGAAAAGGC ACAGCAGGAA CCACTCCGCA GTCTGACAGT 300  
 CAGAACATGC TGAGTGAGAT GGGCAACAAC GGGCTGGATC AGGCCATCAC GCCCGATGGC 360  
 CAGGGCGGCG GGCAGATCGG CGATAATCCT TTAAGTAAAG CCATGCTGAA GCTTATTGCA 420  
 CGCATGATGG ACGGCCAAAG CGATCAGTTT GGCCAACCTG GTACGGGCAA CAACAGTGCC 480  
 TCTTCCGGTA CTTCTTCATC TGGCGGTTCC CCTTTTAAAG ATCTATCAGG GGGGAAGGCC 540  
 CCTTCCGGCA ACTCCCCTTC CGGCAACTAC TCTCCCGTCA GTACCTTCTC ACCCCCATCC 600  
 ACGCCAACGT CCCCTACCTC ACCGCTTGAT TTCCCTTCTT CTCCACCAA AGCAGCCGGG 660  
 GGCAGCACGC CGGTAACCGA TCATCCTGAC CCGTGTGGTA GCGCGGGCAT CGGGGCCGGA 720  
 AATTCGGTGG CCTTCACCAG CGCCGGCGCT AATCAGACGG TGCTGCATGA CACCATTACC 780  
 GTGAAAGCGG GTCAGGTGTT TGATGGCAA GGACAAACCT TCACCGCCGG TTCAGAATTA 840  
 GCGGATGGCG GCCAGTCTGA AAACCAGAAA CCGCTGTTTA TACTGGAAGA CCGTGCCAGC 900  
 CTGAAAAACG TCACCATGGG CGACGACGGG GCGGATGGTA TTCATCTTTA CCGTGATGCC 960  
 AAAATAGACA ATCTGCACGT CACCAACGTG GGTGAGGACG CGATTACCGT TAAGCCAAAC 1020

AGCGCGGGCA AAAAATCCCA CGTTGAAATC ACTAACAGTT CCTTCGAGCA CGCCTCTGAC 1080  
 AAGATCCTGC AGCTGAATGC CGATACTAAC CTGAGCGTTG ACAACGTGAA GGCCAAAGAC 1140  
 TTTGGTACTT TTGTACGCAC TAACGGCGGT CAACAGGGTA ACTGGGATCT GAATCTGAGC 1200  
 CATATCAGCG CAGAAGACGG TAAGTTCTCG TTCGTTAAAA GCGATAGCGA GGGGCTAAAC 1260  
 GTCAATACCA GTGATATCTC ACTGGGTGAT GTTGAAAACC ACTACAAAGT GCCGATGTCC 1320  
 GCCAACCTGA AGGTGGCTGA ATGA 1344

See GenBank Accession No. U94513. The isolated DNA molecule of the present invention encodes a hypersensitive response elicitor protein or polypeptide having an amino acid sequence of SEQ ID NO: 26 as follows:

Met Ser Ile Leu Thr Leu Asn Asn Asn Thr Ser Ser Ser Pro Gly Leu  
 1 5 10 15  
 Phe Gln Ser Gly Gly Asp Asn Gly Leu Gly Gly His Asn Ala Asn Ser  
 20 25 30  
 Ala Leu Gly Gln Gln Pro Ile Asp Arg Gln Thr Ile Glu Gln Met Ala  
 35 40 45  
 Gln Leu Leu Ala Glu Leu Leu Lys Ser Leu Leu Ser Pro Gln Ser Gly  
 50 55 60  
 Asn Ala Ala Thr Gly Ala Gly Gly Asn Asp Gln Thr Thr Gly Val Gly  
 65 70 75 80  
 Asn Ala Gly Gly Leu Asn Gly Arg Lys Gly Thr Ala Gly Thr Thr Pro  
 85 90 95  
 Gln Ser Asp Ser Gln Asn Met Leu Ser Glu Met Gly Asn Asn Gly Leu  
 100 105 110  
 Asp Gln Ala Ile Thr Pro Asp Gly Gln Gly Gly Gly Gln Ile Gly Asp  
 115 120 125  
 Asn Pro Leu Leu Lys Ala Met Leu Lys Leu Ile Ala Arg Met Met Asp  
 130 135 140  
 Gly Gln Ser Asp Gln Phe Gly Gln Pro Gly Thr Gly Asn Asn Ser Ala  
 145 150 155 160  
 Ser Ser Gly Thr Ser Ser Ser Gly Gly Ser Pro Phe Asn Asp Leu Ser  
 165 170 175  
 Gly Gly Lys Ala Pro Ser Gly Asn Ser Pro Ser Gly Asn Tyr Ser Pro  
 180 185 190  
 Val Ser Thr Phe Ser Pro Pro Ser Thr Pro Thr Ser Pro Thr Ser Pro  
 195 200 205  
 Leu Asp Phe Pro Ser Ser Pro Thr Lys Ala Ala Gly Gly Ser Thr Pro  
 210 215 220  
 Val Thr Asp His Pro Asp Pro Val Gly Ser Ala Gly Ile Gly Ala Gly

225					230					235					240
Asn	Ser	Val	Ala	Phe 245	Thr	Ser	Ala	Gly	Ala 250	Asn	Gln	Thr	Val	Leu 255	His
Asp	Thr	Ile	Thr 260	Val	Lys	Ala	Gly	Gln 265	Val	Phe	Asp	Gly	Lys 270	Gly	Gln
Thr	Phe	Thr 275	Ala	Gly	Ser	Glu	Leu 280	Gly	Asp	Gly	Gly	Gln 285	Ser	Glu	Asn
Gln	Lys 290	Pro	Leu	Phe	Ile	Leu 295	Glu	Asp	Gly	Ala	Ser 300	Leu	Lys	Asn	Val
Thr 305	Met	Gly	Asp	Asp	Gly 310	Ala	Asp	Gly	Ile	His 315	Leu	Tyr	Gly	Asp	Ala 320
Lys	Ile	Asp	Asn	Leu 325	His	Val	Thr	Asn	Val 330	Gly	Glu	Asp	Ala	Ile 335	Thr
Val	Lys	Pro	Asn 340	Ser	Ala	Gly	Lys	Lys 345	Ser	His	Val	Glu	Ile 350	Thr	Asn
Ser	Ser	Phe 355	Glu	His	Ala	Ser	Asp 360	Lys	Ile	Leu	Gln	Leu 365	Asn	Ala	Asp
Thr	Asn 370	Leu	Ser	Val	Asp	Asn 375	Val	Lys	Ala	Lys	Asp 380	Phe	Gly	Thr	Phe
Val 385	Arg	Thr	Asn	Gly	Gly 390	Gln	Gln	Gly	Asn	Trp 395	Asp	Leu	Asn	Leu	Ser 400
His	Ile	Ser	Ala	Glu 405	Asp	Gly	Lys	Phe	Ser 410	Phe	Val	Lys	Ser	Asp 415	Ser
Glu	Gly	Leu	Asn 420	Val	Asn	Thr	Ser	Asp 425	Ile	Ser	Leu	Gly	Asp 430	Val	Glu
Asn	His 435	Tyr	Lys	Val	Pro	Met	Ser 440	Ala	Asn	Leu	Lys	Val 445	Ala	Glu	

This protein or polypeptide is acidic, rich in glycine and serine, and lacks cysteine. It is also heat stable, protease sensitive, and suppressed by inhibitors of plant metabolism. The protein or polypeptide of the present invention has a predicted molecular size of ca. 4.5 kDa.



Please replace the paragraph beginning on page 13, line 48, and ending on page 22, line 24, with the following replacement paragraph:

AS  
Another potentially suitable hypersensitive response elicitor from *Erwinia amylovora* is disclosed in U.S. Patent Application Serial No. 09/120,663 which is hereby incorporated by reference. The protein is encoded by a DNA molecule having a nucleic acid sequence of SEQ ID NO: 27 as follows:

```

ATGGAATTAA AATCACTGGG AACTGAACAC AAGCGGCAG TACACACAGC GGCGCACAAAC      60
CCTGTGGGGC ATGGTGTTCG CTTACAGCAG GGCAGCAGCA GCAGCAGCCC GCAAAATGCC      120
GCTGCATCAT TGGCGGCAGA AGGCAAAAAT CGTGGGAAAA TGCCGAGAAT TCACCAGCCA      180
TCTACTGCGG CTGATGGTAT CAGCGCTGCT CACCAGCAAA AGAAATCCTT CAGTCTCAGG      240
GGCTGTTTGG GGACGAAAAA ATTTTCCAGA TCGGCACCGC AGGGCCAGCC AGGTACCACC      300
CACAGCAAAG GGGCAACATT GCGCGATCTG CTGGCGCGGG ACGACGGCGA AACGCAGCAT      360
GAGGCGGCCG CGCCAGATGC GGCGCGTTTG ACCCGTTTCG GCGGCGTCAA ACGCCGCAAT      420
ATGGACGACA TGGCCGGGCG GCCAATGGTG AAAGGTGGCA GCGGCGAAGA TAAGGTACCA      480
ACGCAGCAAA AACGGCATCA GCTGAACAAT TTTGGCCAGA TGCGCCAAAC GATGTTGAGC      540
AAAATGGCTC ACCCGGCTTC AGCCAACGCC GGCATCGCC TGCAGCATTC ACCGCCGCAC      600
ATCCCGGGTA GCCACCACGA AATCAAGGAA GAACCGGTTG GCTCCACCAG CAAGGCAACA      660
ACGCCCCACG CAGACAGAGT GGAAATCGCT CAGGAAGATG ACGACAGCGA ATTCCAGCAA      720
CTGCATCAAC AGCGGCTGGC GCGCGAACGG GAAAATCCAC CGCAGCCGCC CAAACTCGGC      780
GTTGCCACAC CGATTAGCGC CAGGTTTCAG CCCAACTGA CTGCGGTTGC GGAAAGCGTC      840
CTTGAGGGGA CAGATACCAC GCAGTCACCC CTTAAGCCGC AATCAATGCT GAAAGGAAGT      900
GGAGCCGGGG TAACGCCGCT GGCGGTAACG CTGGATAAAG GCAAGTTGCA GCTGGCACCG      960
GATAATCCAC CCGCGCTCAA TACGTTGTTG AAGCAGACAT TGGGTAAAGA CACCCAGCAC     1020
TATCTGGCGC ACCATGCCAG CAGCGACGGT AGCCAGCATC TGCTGCTGGA CAACAAAGGC     1080
CACCTGTTTG ATATCAAAAG CACCGCCACC AGCTATAGCG TGCTGCACAA CAGCCACCCC     1140
GGTGAGATAA AGGGCAAGCT GGCGCAGGCG GGTACTGGCT CCGTCAGCGT AGACGGTAAA     1200
AGCGGCAAGA TCTCGTGGG GAGCGGTACG CAAAGTCACA ACAAACAAT GCTAAGCCAA     1260
CCGGGGGAAG CGCACCGTTC CTTATTAACC GGCATTTGGC AGCATCCTGC TGGCGCAGCG     1320
CGGCCGCAGG GCGAGTCAAT CCGCCTGCAT GACGACAAAA TTCATATCCT GCATCCGGAG     1380
CTGGGCGTAT GGCAATCTGC GGATAAAGAT ACCCACAGCC AGCTGTCTCG CCAGGCAGAC     1440
GGTAAGCTCT ATGCGCTGAA AGACAACCGT ACCCTGCAAA ACCTCTCCGA TAATAAATCC     1500
TCAGAAAAGC TGGTCGATAA AATCAAATCG TATTCCGTTG ATCAGCGGGG GCAGGTGGCG     1560
ATCCTGACGG ATACTCCCGG CCGCCATAAG ATGAGTATTA TGCCCTCGCT GGATGCTTCC     1620

```

CCGGAGAGCC ATATTTCCCT CAGCCTGCAT TTTGCCGATG CCCACCAGGG GTTATTGCAC 1680  
 GGGAAGTCGG AGCTTGAGGC ACAATCTGTC GCGATCAGCC ATGGGCGACT GGTGTGGGCC 1740  
 GATAGCGAAG GCAAGCTGTT TAGCGCCGCC ATTCCGAAGC AAGGGGATGG AAACGAACTG 1800  
 AAAATGAAAG CCATGCCTCA GCATGCGCTC GATGAACATT TTGGTCATGA CCACCAGATT 1860  
 TCTGGATTTT TCCATGACGA CCACGGCCAG CTTAATGCGC TGGTGAAAAA TAACTTCAGG 1920  
 CAGCAGCATG CCTGCCCCTT GGGTAACGAT CATCAGTTTC ACCCCGGCTG GAACCTGACT 1980  
 GATGCGCTGG TTATCGACAA TCAGCTGGGG CTGCATCATA CCAATCCTGA ACCGCATGAG 2040  
 ATTCTTGATA TGGGGCATT T AGGCAGCCTG GCGTTACAGG AGGGCAAGCT TCACTATTTT 2100  
 GACCAGCTGA CCAAAGGGTG GACTGGCGCG GAGTCAGATT GTAAGCAGCT GAAAAAGGC 2160  
 CTGGATGGAG CAGCTTATCT ACTGAAAGAC GGTGAAGTGA AACGCCTGAA TATTAATCAG 2220  
 AGCACCTCCT CTATCAAGCA CGGAACGGAA AACGTTTTTT CGCTGCCGCA TGTGCGCAAT 2280  
 AAACCGGAGC CGGGAGATGC CCTGCAAGGG CTGAATAAAG ACGATAAGGC CCAGGCCATG 2340  
 GCGGTGATTG GGGTAAATAA ATACCTGGCG CTGACGGAAG AAGGGGACAT TCGCTCCTTC 2400  
 CAGATAAAAC CCGGCACCCA GCAGTTGGAG CGGCCGGCAC AAATCTCAG CCGCGAAGGT 2460  
 ATCAGCGGCG AACTGAAAGA CATTCATGTC GACCACAAGC AGAACCTGTA TGCCTTGACC 2520  
 CACGAGGGAG AGGTGTTTCA TCAGCCGCGT GAAGCCTGGC AGAATGGTGC CGAAAGCAGC 2580  
 AGCTGGCACA AACTGGCGTT GCCACAGAGT GAAAGTAAGC TAAAAAGTCT GGACATGAGC 2640  
 CATGAGCACA AACCGATTGC CACCTTTGAA GACGGTAGCC AGCATCAGCT GAAGGCTGGC 2700  
 GGCTGGCACG CCTATGCGGC ACCTGAACGC GGGCCGCTGG CCGTGGGTAC CAGCGGTTCA 2760  
 CAAACCGTCT TTAACCGACT AATGCAGGGG GTGAAAGGCA AGGTGATCCC AGGCAGCGGG 2820  
 TTGACGGTTA AGCTCTCGGC TCAGACGGGG GGAATGACCG GCGCCGAAGG GCGCAAGGTC 2880  
 AGCAGTAAAT TTTCCGAAAG GATCCGCGCC TATGCGTTCA ACCCAACAAT GTCCACGCCG 2940  
 CGACCGATTA AAAATGCTGC TTATGCCACA CAGCACGGCT GGCAGGGGCG TGAGGGGTTG 3000  
 AAGCCGTTGT ACGAGATGCA GGGAGCGCTG ATTAAACAAC TGGATGCGCA TAACGTTCTG 3060  
 CATAACGCGC CACAGCCAGA TTTGCAGAGC AAATGGAAG CTCTGGATTT AGGCGAACAT 3120  
 GGCGCAGAAT TGCTTAACGA CATGAAGCGC TTCCGCGACG AACTGGAGCA GAGTGCAACC 3180  
 CGTTCGGTGA CCGTTTTAGG TCAACATCAG GGAGTGCTAA AAAGCAACGG TGAAATCAAT 3240  
 AGCGAATTTA AGCCATCGCC CGGCAAGGCG TTGGTCCAGA GCTTTAACGT CAATCGCTCT 3300  
 GGTGAGGATC TAAGCAAGTC ACTGCAACAG GCAGTACATG CCACGCCGCC ATCCGCAGAG 3360  
 AGTAACTGC AATCCATGCT GGGGCACTTT GTCACTGCGG GGGTGGATAT GAGTCATCAG 3420  
 AAGGGCGAGA TCCCGCTGGG CCGCCAGCGC GATCCGAATG ATAAAACCGC ACTGACCAAA 3480  
 TCGCGTTTAA TTTTAGATAC CGTGACCATC GGTGAACTGC ATGAACTGGC CGATAAGGCG 3540  
 AAATGGTAT CTGACCATAA ACCCGATGCC GATCAGATAA AACAGCTGCG CCAGCAGTTC 3600

GATACGCTGC GTGAAAAGCG GTATGAGAGC AATCCGGTGA AGCATTACAC CGATATGGGC 3660  
 TTCACCCATA ATAAGGCGCT GGAAGCAAAC TATGATGCGG TCAAAGCCTT TATCAATGCC 3720  
 TTTAAGAAAG AGCACCACGG CGTCAATCTG ACCACGCGTA CCGTACTGGA ATCACAGGGC 3780  
 AGTGCGGAGC TGGCGAAGAA GCTCAAGAAT ACGCTGTTGT CCCTGGACAG TGGTGAAAGT 3840  
 ATGAGCTTCA GCCGGTCATA TGGCGGGGGC GTCAGCACTG TCTTTGTGCC TACCCTTAGC 3900  
 AAGAAGGTGC CAGTTCCGGT GATCCCCGGA GCCGGCATCA CGCTGGATCG CGCCTATAAC 3960  
 CTGAGCTTCA GTCGTACCAG CGGCGGATTG AACGTCAGTT TTGGCCGCGA CGGCGGGGTG 4020  
 AGTGTAACA TCATGGTCGC TACCGGCCAT GATGTGATGC CCTATATGAC CGGTAAGAAA 4080  
 ACCAGTGCAG GTAACGCCAG TGA CTGGTTG AGCGCAAAAC ATAAAATCAG CCCGGACTTG 4140  
 CGTATCGGCG CTGCTGTGAG TGGCACCTTG CAAGGAACGC TACAAAACAG CCTGAAGTTT 4200  
 AAGCTGACAG AGGATGAGCT GCCTGGCTTT ATCCATGGCT TGACGCATGG CACGTTGACC 4260  
 CCGGCAGAAC TGTTGCAAAA GGGGATCGAA CATCAGATGA AGCAGGGCAG CAAACTGACG 4320  
 TTTAGCGTCG ATACCTCGGC AAATCTGGAT CTGCGTGCCG GTATCAATCT GAACGAAGAC 4380  
 GGCAGTAAAC CAAATGGTGT CACTGCCCCT GTTCTGCCC GGCTAAGTGC ATCGGCAAAC 4440  
 CTGGCCGCCG GCTCGCGTGA ACGCAGCACC ACCTCTGGCC AGTTTGGCAG CACGACTTCG 4500  
 GCCAGCAATA ACCGCCCAAC CTTCTCAAC GGGGTGCGCG CGGGTGCTAA CCTGACGGCT 4560  
 GCTTTAGGGG TTGCCATTTC ATCTACGCAT GAAGGGAAAC CGGTCGGGAT CTTCCCGGCA 4620  
 TTTACCTCGA CCAATGTTTC GGCAGCGCTG GCGCTGGATA ACCGTACCTC ACAGAGTATC 4680  
 AGCCTGGAAT TGAAGCGCGC GGAGCCGGTG ACCAGCAACG ATATCAGCGA GTTGACCTCC 4740  
 ACGCTGGGAA AACACTTTAA GGATAGCGCC ACAACGAAGA TGCTTGCCGC TCTCAAAGAG 4800  
 TTAGATGACG CTAAGCCCGC TGAACAACTG CATATTTTAC AGCAGCATTT CAGTGCAAAA 4860  
 GATGTCGTCG GTGATGAACG CTACGAGGCG GTGCGCAACC TGAAAAAACT GGTGATACGT 4920  
 CAACAGGCTG CGGACAGCCA CAGCATGGAA TTAGGATCTG CCAGTCACAG CACGACCTAC 4980  
 AATAATCTGT CGAGAATAAA TAATGACGGC ATTGTCGAGC TGCTACACAA ACATTTCGAT 5040  
 GCGGCATTAC CAGCAAGCAG TGCCAAACGT CTTGGTGAAA TGATGAATAA CGATCCGGCA 5100  
 CTGAAAGATA TTATTAAGCA GCTGCAAAGT ACGCCGTTCA GCAGCGCCAG CGTGTCGATG 5160  
 GAGCTGAAAG ATGGTCTGCG TGAGCAGACG GAAAAAGCAA TACTGGACGG TAAGGTCGGT 5220  
 CGTGAAGAAG TGGGAGTACT TTTCCAGGAT CGTAACAACT TGCGTGTTAA ATCGGTCAGC 5280  
 GTCAGTCAGT CCGTCAGCAA AAGCGAAGGC TTCAATACCC CAGCGCTGTT ACTGGGGACG 5340  
 AGCAACAGCG CTGCTATGAG CATGGAGCGC AACATCGGAA CCATTAATTT TAAATACGGC 5400  
 CAGGATCAGA ACACCCACG GCGATTTACC CTGGAGGGTG GAATAGCTCA GGCTAATCCG 5460  
 CAGGTCGCAT CTGCGCTTAC TGATTTGAAG AAGGAAGGGC TGGAAATGAA GAGCTAA 5517

AS  
Gm  
This DNA molecule is known as the dspE gene for *Erwinia amylovora*. This isolated DNA molecule of the present invention encodes a protein or polypeptide which elicits a plant pathogen's hypersensitive response having an amino acid sequence of SEQ ID NO: 28 as follows:

Met	Glu	Leu	Lys	Ser	Leu	Gly	Thr	Glu	His	Lys	Ala	Ala	Val	His	Thr	1	5	10	15
Ala	Ala	His	Asn	Pro	Val	Gly	His	Gly	Val	Ala	Leu	Gln	Gln	Gly	Ser	20	25	30	
Ser	Ser	Ser	Ser	Pro	Gln	Asn	Ala	Ala	Ala	Ser	Leu	Ala	Ala	Glu	Gly	35	40	45	
Lys	Asn	Arg	Gly	Lys	Met	Pro	Arg	Ile	His	Gln	Pro	Ser	Thr	Ala	Ala	50	55	60	
Asp	Gly	Ile	Ser	Ala	Ala	His	Gln	Gln	Lys	Lys	Ser	Phe	Ser	Leu	Arg	65	70	75	80
Gly	Cys	Leu	Gly	Thr	Lys	Lys	Phe	Ser	Arg	Ser	Ala	Pro	Gln	Gly	Gln	85	90	95	
Pro	Gly	Thr	Thr	His	Ser	Lys	Gly	Ala	Thr	Leu	Arg	Asp	Leu	Leu	Ala	100	105	110	
Arg	Asp	Asp	Gly	Glu	Thr	Gln	His	Glu	Ala	Ala	Ala	Pro	Asp	Ala	Ala	115	120	125	
Arg	Leu	Thr	Arg	Ser	Gly	Gly	Val	Lys	Arg	Arg	Asn	Met	Asp	Asp	Met	130	135	140	
Ala	Gly	Arg	Pro	Met	Val	Lys	Gly	Gly	Ser	Gly	Glu	Asp	Lys	Val	Pro	145	150	155	160
Thr	Gln	Gln	Lys	Arg	His	Gln	Leu	Asn	Asn	Phe	Gly	Gln	Met	Arg	Gln	165	170	175	
Thr	Met	Leu	Ser	Lys	Met	Ala	His	Pro	Ala	Ser	Ala	Asn	Ala	Gly	Asp	180	185	190	
Arg	Leu	Gln	His	Ser	Pro	Pro	His	Ile	Pro	Gly	Ser	His	His	Glu	Ile	195	200	205	
Lys	Glu	Glu	Pro	Val	Gly	Ser	Thr	Ser	Lys	Ala	Thr	Thr	Ala	His	Ala	210	215	220	
Asp	Arg	Val	Glu	Ile	Ala	Gln	Glu	Asp	Asp	Asp	Ser	Glu	Phe	Gln	Gln	225	230	235	240
Leu	His	Gln	Gln	Arg	Leu	Ala	Arg	Glu	Arg	Glu	Asn	Pro	Pro	Gln	Pro	245	250	255	
Pro	Lys	Leu	Gly	Val	Ala	Thr	Pro	Ile	Ser	Ala	Arg	Phe	Gln	Pro	Lys	260	265	270	
Leu	Thr	Ala	Val	Ala	Glu	Ser	Val	Leu	Glu	Gly	Thr	Asp	Thr	Thr	Gln	275	280	285	
Ser	Pro	Leu	Lys	Pro	Gln	Ser	Met	Leu	Lys	Gly	Ser	Gly	Ala	Gly	Val	290	295	300	
Thr	Pro	Leu	Ala	Val	Thr	Leu	Asp	Lys	Gly	Lys	Leu	Gln	Leu	Ala	Pro	305	310	315	320

Asp Asn Pro Pro Ala Leu Asn Thr Leu Leu Lys Gln Thr Leu Gly Lys  
 325 330 335  
 Asp Thr Gln His Tyr Leu Ala His His Ala Ser Ser Asp Gly Ser Gln  
 340 345 350  
 His Leu Leu Leu Asp Asn Lys Gly His Leu Phe Asp Ile Lys Ser Thr  
 355 360 365  
 Ala Thr Ser Tyr Ser Val Leu His Asn Ser His Pro Gly Glu Ile Lys  
 370 375 380  
 Gly Lys Leu Ala Gln Ala Gly Thr Gly Ser Val Ser Val Asp Gly Lys  
 385 390 395 400  
 Ser Gly Lys Ile Ser Leu Gly Ser Gly Thr Gln Ser His Asn Lys Thr  
 405 410 415  
 Met Leu Ser Gln Pro Gly Glu Ala His Arg Ser Leu Leu Thr Gly Ile  
 420 425 430  
 Trp Gln His Pro Ala Gly Ala Ala Arg Pro Gln Gly Glu Ser Ile Arg  
 435 440 445  
 Leu His Asp Asp Lys Ile His Ile Leu His Pro Glu Leu Gly Val Trp  
 450 455 460  
 Gln Ser Ala Asp Lys Asp Thr His Ser Gln Leu Ser Arg Gln Ala Asp  
 465 470 475 480  
 Gly Lys Leu Tyr Ala Leu Lys Asp Asn Arg Thr Leu Gln Asn Leu Ser  
 485 490 495  
 Asp Asn Lys Ser Ser Glu Lys Leu Val Asp Lys Ile Lys Ser Tyr Ser  
 500 505 510  
 Val Asp Gln Arg Gly Gln Val Ala Ile Leu Thr Asp Thr Pro Gly Arg  
 515 520 525  
 His Lys Met Ser Ile Met Pro Ser Leu Asp Ala Ser Pro Glu Ser His  
 530 535 540  
 Ile Ser Leu Ser Leu His Phe Ala Asp Ala His Gln Gly Leu Leu His  
 545 550 555 560  
 Gly Lys Ser Glu Leu Glu Ala Gln Ser Val Ala Ile Ser His Gly Arg  
 565 570 575  
 Leu Val Val Ala Asp Ser Glu Gly Lys Leu Phe Ser Ala Ala Ile Pro  
 580 585 590  
 Lys Gln Gly Asp Gly Asn Glu Leu Lys Met Lys Ala Met Pro Gln His  
 595 600 605  
 Ala Leu Asp Glu His Phe Gly His Asp His Gln Ile Ser Gly Phe Phe  
 610 615 620  
 His Asp Asp His Gly Gln Leu Asn Ala Leu Val Lys Asn Asn Phe Arg  
 625 630 635 640  
 Gln Gln His Ala Cys Pro Leu Gly Asn Asp His Gln Phe His Pro Gly  
 645 650 655  
 Trp Asn Leu Thr Asp Ala Leu Val Ile Asp Asn Gln Leu Gly Leu His  
 660 665 670

His Thr Asn Pro Glu Pro His Glu Ile Leu Asp Met Gly His Leu Gly  
 675 680 685  
 Ser Leu Ala Leu Gln Glu Gly Lys Leu His Tyr Phe Asp Gln Leu Thr  
 690 695 700  
 Lys Gly Trp Thr Gly Ala Glu Ser Asp Cys Lys Gln Leu Lys Lys Gly  
 705 710 715 720  
 Leu Asp Gly Ala Ala Tyr Leu Leu Lys Asp Gly Glu Val Lys Arg Leu  
 725 730 735  
 Asn Ile Asn Gln Ser Thr Ser Ser Ile Lys His Gly Thr Glu Asn Val  
 740 745 750  
 Phe Ser Leu Pro His Val Arg Asn Lys Pro Glu Pro Gly Asp Ala Leu  
 755 760 765  
 Gln Gly Leu Asn Lys Asp Asp Lys Ala Gln Ala Met Ala Val Ile Gly  
 770 775 780  
 Val Asn Lys Tyr Leu Ala Leu Thr Glu Lys Gly Asp Ile Arg Ser Phe  
 785 790 795 800  
 Gln Ile Lys Pro Gly Thr Gln Gln Leu Glu Arg Pro Ala Gln Thr Leu  
 805 810 815  
 Ser Arg Glu Gly Ile Ser Gly Glu Leu Lys Asp Ile His Val Asp His  
 820 825 830  
 Lys Gln Asn Leu Tyr Ala Leu Thr His Glu Gly Glu Val Phe His Gln  
 835 840 845  
 Pro Arg Glu Ala Trp Gln Asn Gly Ala Glu Ser Ser Ser Trp His Lys  
 850 855 860  
 Leu Ala Leu Pro Gln Ser Glu Ser Lys Leu Lys Ser Leu Asp Met Ser  
 865 870 875 880  
 His Glu His Lys Pro Ile Ala Thr Phe Glu Asp Gly Ser Gln His Gln  
 885 890 895  
 Leu Lys Ala Gly Gly Trp His Ala Tyr Ala Ala Pro Glu Arg Gly Pro  
 900 905 910  
 Leu Ala Val Gly Thr Ser Gly Ser Gln Thr Val Phe Asn Arg Leu Met  
 915 920 925  
 Gln Gly Val Lys Gly Lys Val Ile Pro Gly Ser Gly Leu Thr Val Lys  
 930 935 940  
 Leu Ser Ala Gln Thr Gly Gly Met Thr Gly Ala Glu Gly Arg Lys Val  
 945 950 955 960  
 Ser Ser Lys Phe Ser Glu Arg Ile Arg Ala Tyr Ala Phe Asn Pro Thr  
 965 970 975  
 Met Ser Thr Pro Arg Pro Ile Lys Asn Ala Ala Tyr Ala Thr Gln His  
 980 985 990  
 Gly Trp Gln Gly Arg Glu Gly Leu Lys Pro Leu Tyr Glu Met Gln Gly  
 995 1000 1005  
 Ala Leu Ile Lys Gln Leu Asp Ala His Asn Val Arg His Asn Ala Pro  
 1010 1015 1020

P5  
 Gln Pro Asp Leu Gln Ser Lys Leu Glu Thr Leu Asp Leu Gly Glu His  
 1025 1030 1035 1040  
 Gly Ala Glu Leu Leu Asn Asp Met Lys Arg Phe Arg Asp Glu Leu Glu  
 1045 1050 1055  
 Gln Ser Ala Thr Arg Ser Val Thr Val Leu Gly Gln His Gln Gly Val  
 1060 1065 1070  
 Leu Lys Ser Asn Gly Glu Ile Asn Ser Glu Phe Lys Pro Ser Pro Gly  
 1075 1080 1085  
 Lys Ala Leu Val Gln Ser Phe Asn Val Asn Arg Ser Gly Gln Asp Leu  
 1090 1095 1100  
 Ser Lys Ser Leu Gln Gln Ala Val His Ala Thr Pro Pro Ser Ala Glu  
 1105 1110 1115 1120  
 Ser Lys Leu Gln Ser Met Leu Gly His Phe Val Ser Ala Gly Val Asp  
 1125 1130 1135  
 Met Ser His Gln Lys Gly Glu Ile Pro Leu Gly Arg Gln Arg Asp Pro  
 1140 1145 1150  
 Asn Asp Lys Thr Ala Leu Thr Lys Ser Arg Leu Ile Leu Asp Thr Val  
 1155 1160 1165  
 Thr Ile Gly Glu Leu His Glu Leu Ala Asp Lys Ala Lys Leu Val Ser  
 1170 1175 1180  
 Asp His Lys Pro Asp Ala Asp Gln Ile Lys Gln Leu Arg Gln Gln Phe  
 1185 1190 1195 1200  
 Asp Thr Leu Arg Glu Lys Arg Tyr Glu Ser Asn Pro Val Lys His Tyr  
 1205 1210 1215  
 Thr Asp Met Gly Phe Thr His Asn Lys Ala Leu Glu Ala Asn Tyr Asp  
 1220 1225 1230  
 Ala Val Lys Ala Phe Ile Asn Ala Phe Lys Lys Glu His His Gly Val  
 1235 1240 1245  
 Asn Leu Thr Thr Arg Thr Val Leu Glu Ser Gln Gly Ser Ala Glu Leu  
 1250 1255 1260  
 Ala Lys Lys Leu Lys Asn Thr Leu Leu Ser Leu Asp Ser Gly Glu Ser  
 1265 1270 1275 1280  
 Met Ser Phe Ser Arg Ser Tyr Gly Gly Gly Val Ser Thr Val Phe Val  
 1285 1290 1295  
 Pro Thr Leu Ser Lys Lys Val Pro Val Pro Val Ile Pro Gly Ala Gly  
 1300 1305 1310  
 Ile Thr Leu Asp Arg Ala Tyr Asn Leu Ser Phe Ser Arg Thr Ser Gly  
 1315 1320 1325  
 Gly Leu Asn Val Ser Phe Gly Arg Asp Gly Gly Val Ser Gly Asn Ile  
 1330 1335 1340  
 Met Val Ala Thr Gly His Asp Val Met Pro Tyr Met Thr Gly Lys Lys  
 1345 1350 1355 1360  
 Thr Ser Ala Gly Asn Ala Ser Asp Trp Leu Ser Ala Lys His Lys Ile  
 1365 1370 1375

Ser Pro Asp Leu Arg Ile Gly Ala Ala Val Ser Gly Thr Leu Gln Gly  
 1380 1385 1390  
 Thr Leu Gln Asn Ser Leu Lys Phe Lys Leu Thr Glu Asp Glu Leu Pro  
 1395 1400 1405  
 Gly Phe Ile His Gly Leu Thr His Gly Thr Leu Thr Pro Ala Glu Leu  
 1410 1415 1420  
 Leu Gln Lys Gly Ile Glu His Gln Met Lys Gln Gly Ser Lys Leu Thr  
 1425 1430 1435 1440  
 Phe Ser Val Asp Thr Ser Ala Asn Leu Asp Leu Arg Ala Gly Ile Asn  
 1445 1450 1455  
 Leu Asn Glu Asp Gly Ser Lys Pro Asn Gly Val Thr Ala Arg Val Ser  
 1460 1465 1470  
 Ala Gly Leu Ser Ala Ser Ala Asn Leu Ala Ala Gly Ser Arg Glu Arg  
 1475 1480 1485  
 Ser Thr Thr Ser Gly Gln Phe Gly Ser Thr Thr Ser Ala Ser Asn Asn  
 1490 1495 1500  
 Arg Pro Thr Phe Leu Asn Gly Val Gly Ala Gly Ala Asn Leu Thr Ala  
 1505 1510 1515 1520  
 Ala Leu Gly Val Ala His Ser Ser Thr His Glu Gly Lys Pro Val Gly  
 1525 1530 1535  
 Ile Phe Pro Ala Phe Thr Ser Thr Asn Val Ser Ala Ala Leu Ala Leu  
 1540 1545 1550  
 Asp Asn Arg Thr Ser Gln Ser Ile Ser Leu Glu Leu Lys Arg Ala Glu  
 1555 1560 1565  
 Pro Val Thr Ser Asn Asp Ile Ser Glu Leu Thr Ser Thr Leu Gly Lys  
 1570 1575 1580  
 His Phe Lys Asp Ser Ala Thr Thr Lys Met Leu Ala Ala Leu Lys Glu  
 1585 1590 1595 1600  
 Leu Asp Asp Ala Lys Pro Ala Glu Gln Leu His Ile Leu Gln Gln His  
 1605 1610 1615  
 Phe Ser Ala Lys Asp Val Val Gly Asp Glu Arg Tyr Glu Ala Val Arg  
 1620 1625 1630  
 Asn Leu Lys Lys Leu Val Ile Arg Gln Gln Ala Ala Asp Ser His Ser  
 1635 1640 1645  
 Met Glu Leu Gly Ser Ala Ser His Ser Thr Thr Tyr Asn Asn Leu Ser  
 1650 1655 1660  
 Arg Ile Asn Asn Asp Gly Ile Val Glu Leu Leu His Lys His Phe Asp  
 1665 1670 1675 1680  
 Ala Ala Leu Pro Ala Ser Ser Ala Lys Arg Leu Gly Glu Met Met Asn  
 1685 1690 1695  
 Asn Asp Pro Ala Leu Lys Asp Ile Ile Lys Gln Leu Gln Ser Thr Pro  
 1700 1705 1710  
 Phe Ser Ser Ala Ser Val Ser Met Glu Leu Lys Asp Gly Leu Arg Glu  
 1715 1720 1725



ASG  
cont'd

Gln Thr Glu Lys Ala Ile Leu Asp Gly Lys Val Gly Arg Glu Glu Val  
1730 1735 1740

Gly Val Leu Phe Gln Asp Arg Asn Asn Leu Arg Val Lys Ser Val Ser  
1745 1750 1755 1760

Val Ser Gln Ser Val Ser Lys Ser Glu Gly Phe Asn Thr Pro Ala Leu  
1765 1770 1775

Leu Leu Gly Thr Ser Asn Ser Ala Ala Met Ser Met Glu Arg Asn Ile  
1780 1785 1790

Gly Thr Ile Asn Phe Lys Tyr Gly Gln Asp Gln Asn Thr Pro Arg Arg  
1795 1800 1805

Phe Thr Leu Glu Gly Gly Ile Ala Gln Ala Asn Pro Gln Val Ala Ser  
1810 1815 1820

Ala Leu Thr Asp Leu Lys Lys Glu Gly Leu Glu Met Lys Ser  
1825 1830 1835

This protein or polypeptide is about 198 kDa and has a pI of 8.98.

Please replace the paragraph beginning on page 22, line 25, and ending on page 23, line 17, with the following replacement paragraph:

AL The present invention relates to an isolated DNA molecule having a nucleotide sequence of SEQ ID NO: 29 as follows:

ATGACATCGT CACAGCAGCG GGTGAAAGG TTTTACAGT ATTTCTCCGC CGGGTGTAAG 60

ACGCCATAC ATCTGAAAGA CGGGGTGTGC GCCCTGTATA ACGAACAAGA TGAGGAGGCG 120

GCGGTGCTGG AAGTACCGCA ACACAGCGAC AGCCTGTTAC TACACTGCCG AATCATTGAG 180

GCTGACCCAC AAACCTCAAT AACCTGTAT TCGATGCTAT TACAGCTGAA TTTTGAAATG 240

GCGGCCATGC GCGGCTGTTG GCTGGCGCTG GATGAACTGC ACAACGTGCG TTTATGTTTT 300

CAGCAGTCGC TGGAGCATCT GGATGAAGCA AGTTTATAGC ATATCGTTAG CGGCTTCATC 360

GAACATGCGG CAGAAGTGCG TGAGTATATA GCGCAATTAG ACGAGAGTAG CGCGGCATAA 420

This is known as the dspF gene. This isolated DNA molecule of the present invention encodes a hypersensitive response elicitor protein or polypeptide having an amino acid sequence of SEQ ID NO: 30 as follows:

Met Thr Ser Ser Gln Gln Arg Val Glu Arg Phe Leu Gln Tyr Phe Ser  
1 5 10 15

Ala Gly Cys Lys Thr Pro Ile His Leu Lys Asp Gly Val Cys Ala Leu  
20 25 30

Tyr Asn Glu Gln Asp Glu Glu Ala Ala Val Leu Glu Val Pro Gln His  
 35 40 45  
 Ser Asp Ser Leu Leu Leu His Cys Arg Ile Ile Glu Ala Asp Pro Gln  
 50 55 60  
 Thr Ser Ile Thr Leu Tyr Ser Met Leu Leu Gln Leu Asn Phe Glu Met  
 65 70 75 80  
 Ala Ala Met Arg Gly Cys Trp Leu Ala Leu Asp Glu Leu His Asn Val  
 85 90 95  
 Arg Leu Cys Phe Gln Gln Ser Leu Glu His Leu Asp Glu Ala Ser Phe  
 100 105 110  
 Ser Asp Ile Val Ser Gly Phe Ile Glu His Ala Ala Glu Val Arg Glu  
 115 120 125  
 Tyr Ile Ala Gln Leu Asp Glu Ser Ser Ala Ala  
 130 135

This protein or polypeptide is about 16 kDa and has a pI of 4.45.

Please replace the paragraph beginning on page 23, line 18, and ending on page 25, line 13, with the following replacement paragraph:

The hypersensitive response elicitor polypeptide or protein derived from *Pseudomonas syringae* has an amino acid sequence corresponding to SEQ ID NO: 31 as follows:

Met Gln Ser Leu Ser Leu Asn Ser Ser Ser Leu Gln Thr Pro Ala Met  
 1 5 10 15  
 Ala Leu Val Leu Val Arg Pro Glu Ala Glu Thr Thr Gly Ser Thr Ser  
 20 25 30  
 Ser Lys Ala Leu Gln Glu Val Val Val Lys Leu Ala Glu Glu Leu Met  
 35 40 45  
 Arg Asn Gly Gln Leu Asp Asp Ser Ser Pro Leu Gly Lys Leu Leu Ala  
 50 55 60  
 Lys Ser Met Ala Ala Asp Gly Lys Ala Gly Gly Gly Ile Glu Asp Val  
 65 70 75 80  
 Ile Ala Ala Leu Asp Lys Leu Ile His Glu Lys Leu Gly Asp Asn Phe  
 85 90 95  
 Gly Ala Ser Ala Asp Ser Ala Ser Gly Thr Gly Gln Gln Asp Leu Met  
 100 105 110  
 Thr Gln Val Leu Asn Gly Leu Ala Lys Ser Met Leu Asp Asp Leu Leu  
 115 120 125

A7  
 Cont'd

Thr	Lys	Gln	Asp	Gly	Gly	Thr	Ser	Phe	Ser	Glu	Asp	Asp	Met	Pro	Met	
130						135					140					
Leu	Asn	Lys	Ile	Ala	Gln	Phe	Met	Asp	Asp	Asn	Pro	Ala	Gln	Phe	Pro	
145					150					155					160	
Lys	Pro	Asp	Ser	Gly	Ser	Trp	Val	Asn	Glu	Leu	Lys	Glu	Asp	Asn	Phe	
				165					170					175		
Leu	Asp	Gly	Asp	Glu	Thr	Ala	Ala	Phe	Arg	Ser	Ala	Leu	Asp	Ile	Ile	
			180					185					190			
Gly	Gln	Gln	Leu	Gly	Asn	Gln	Gln	Ser	Asp	Ala	Gly	Ser	Leu	Ala	Gly	
		195				200						205				
Thr	Gly	Gly	Gly	Leu	Gly	Thr	Pro	Ser	Ser	Phe	Ser	Asn	Asn	Ser	Ser	
210						215					220					
Val	Met	Gly	Asp	Pro	Leu	Ile	Asp	Ala	Asn	Thr	Gly	Pro	Gly	Asp	Ser	
225					230					235					240	
Gly	Asn	Thr	Arg	Gly	Glu	Ala	Gly	Gln	Leu	Ile	Gly	Glu	Leu	Ile	Asp	
				245					250					255		
Arg	Gly	Leu	Gln	Ser	Val	Leu	Ala	Gly	Gly	Gly	Leu	Gly	Thr	Pro	Val	
			260					265					270			
Asn	Thr	Pro	Gln	Thr	Gly	Thr	Ser	Ala	Asn	Gly	Gly	Gln	Ser	Ala	Gln	
		275					280					285				
Asp	Leu	Asp	Gln	Leu	Leu	Gly	Gly	Leu	Leu	Leu	Lys	Gly	Leu	Glu	Ala	
		290				295					300					
Thr	Leu	Lys	Asp	Ala	Gly	Gln	Thr	Gly	Thr	Asp	Val	Gln	Ser	Ser	Ala	
305					310					315					320	
Ala	Gln	Ile	Ala	Thr	Leu	Leu	Val	Ser	Thr	Leu	Leu	Gln	Gly	Thr	Arg	
				325					330					335		
Asn	Gln	Ala	Ala	Ala												
				340												

This hypersensitive response elicitor polypeptide or protein has a molecular weight of 34-35 kDa. It is rich in glycine (about 13.5%) and lacks cysteine and tyrosine. Further information about the hypersensitive response elicitor derived from *Pseudomonas syringae* is found in He, S. Y., H. C. Huang, and A. Collmer, "*Pseudomonas syringae* pv. *syringae* Harpin<sub>PSS</sub>: a Protein that is Secreted via the Hrp Pathway and Elicits the Hypersensitive Response in Plants," Cell 73:1255-1266 (1993), which is hereby incorporated by reference. The DNA molecule encoding the hypersensitive response elicitor from *Pseudomonas syringae* has a nucleotide sequence corresponding to SEQ ID NO: 32 as follows:

ATGCAGAGTC TCAGTCTTAA CAGCAGCTCG CTGCAAACCC CGGCAATGGC CCTTGTCTCTG

60

GTACGTCCTG AAGCCGAGAC GACTGGCAGT ACGTCGAGCA AGGCGCTTCA GGAAGTTGTC 120  
 GTGAAGCTGG CCGAGGAACT GATGCGCAAT GGTCAACTCG ACGACAGCTC GCCATTGGGA 180  
 AAAGTGTGG CCAAGTCGAT GGCCGCAGAT GGCAAGGCGG GCGGCGGTAT TGAGGATGTC 240  
 ATCGCTGCGC TGGACAAGCT GATCCATGAA AAGCTCGGTG ACAACTTCGG CGCGTCTGCG 300  
 GACAGCGCCT CGGGTACCGG ACAGCAGGAC CTGATGACTC AGGTGCTCAA TGGCCTGGCC 360  
 AAGTCGATGC TCGATGATCT TCTGACCAAG CAGGATGGCG GGACAAGCTT CTCCGAAGAC 420  
 GATATGCCGA TGCTGAACAA GATCGCGCAG TTCATGGATG ACAATCCCGC ACAGTTTCCC 480  
 AAGCCGGACT CGGGCTCCTG GGTGAACGAA CTCAAGGAAG ACAACTTCCT TGATGGCGAC 540  
 GAAACGGCTG CGTTCCGTTT GGCACCTGAC ATCATTGGCC AGCAACTGGG TAATCAGCAG 600  
 AGTGACGCTG GCAGTCTGGC AGGGACGGGT GGAGGTCTGG GCACTCCGAG CAGTTTTTCC 660  
 AACAACTCGT CCGTGATGGG TGATCCGCTG ATCGACGCCA ATACCGGTCC CGGTGACAGC 720  
 GGCAATACCC GTGGTGAAGC GGGGCAACTG ATCGGCAGC TTATCGACCG TGGCCTGCAA 780  
 TCGGTATTGG CCGGTGGTGG ACTGGGCACA CCCGTAAACA CCCCAGAC CCGTACGTCG 840  
 GCGAATGGCG GACAGTCCGC TCAGGATCTT GATCAGTTGC TGGGCGGCTT GCTGCTCAAG 900  
 GGCCTGGAGG CAACGCTCAA GGATGCCGGG CAAACAGGCA CCGACGTGCA GTCGAGCGCT 960  
 GCGCAAATCG CCACCTTGCT GGTGAGTACG CTGCTGCAAG GCACCCGCAA TCAGGCTGCA 1020  
 GCCTGA 1026

Please replace the paragraph beginning on page 25, line 15, and ending on page 28, line 6, with the following replacement paragraph:

Another potentially suitable hypersensitive response elicitor from  
*Pseudomonas syringae* is disclosed in U.S. Patent Application Serial No. 09/120,817, which  
 is hereby incorporated by reference. The protein has a nucleotide sequence of SEQ ID  
 NO: 33 as follows:

TCCACTTCGC TGATTTTGAA ATTGGCAGAT TCATAGAAAC GTTCAGGTGT GGAAATCAGG 60  
 CTGAGTGCGC AGATTTCGTT GATAAGGGTG TGGTACTGGT CATTGTTGGT CATTTCAGG 120  
 CCTCTGAGTG CCGTGCGGAG CAATACCAGT CTTCTGCTG GCGTGTGCAC ACTGAGTCGC 180  
 AGGCATAGGC ATTTTCAGTTC CTTGCGTTGG TTGGGCATAT AAAAAAGGA ACTTTTAAAA 240  
 ACAGTGCAAT GAGATGCCGG CAAAACGGGA ACCGGTCGCT GCGCTTTGCC ACTCACTTCG 300  
 AGCAAGCTCA ACCCAAACA TCCACATCCC TATCGAACGG ACAGCGATAC GGCCACTTGC 360

TCTGGTAAAC CCTGGAGCTG GCGTCGGTCC AATTGCCCAC TTAGCGAGGT AACGCAGCAT 420  
 GAGCATCGGC ATCACACCCC GGCCGCAACA GACCACCACG CCACTCGATT TTTCGGCGCT 480  
 AAGCGGCAAG AGTCCTCAAC CAAACACGTT CGGCGAGCAG AACACTCAGC AAGCGATCGA 540  
 CCCGAGTGCA CTGTTGTTTCG GCAGCGACAC ACAGAAAGAC GTCAACTTCG GCACGCCCGA 600  
 CAGCACCCTC CAGAATCCGC AGGACGCCAG CAAGCCCAAC GACAGCCAGT CCAACATCGC 660  
 TAAATTGATC AGTGATTGA TCATGTCGTT GCTGCAGATG CTCACCAACT CCAATAAAAA 720  
 GCAGGACACC AATCAGGAAC AGCCTGATAG CCAGGCTCCT TTCCAGAACA ACGGCGGGCT 780  
 CGGTACACCG TCGGCCGATA GCGGGGGCGG CGGTACACCG GATGCGACAG GTGGCGGCGG 840  
 CGGTGATACG CCAAGCGCAA CAGGCGGTGG CGGCGGTGAT ACTCCGACCG CAACAGGCGG 900  
 TGGCGGCAGC GGTGGCGGCG GCACACCCAC TGCAACAGGT GGCAGCAGCG GTGGCACACC 960  
 CACTGCAACA GGCGGTGGCG AGGGTGGCGT AACACCGCAA ATCACTCCGC AGTTGGCCAA 1020  
 CCCTAACCGT ACCTCAGGTA CTGGCTCGGT GTCGGACACC GCAGGTTCTA CCGAGCAAGC 1080  
 CGGCAAGATC AATGTGGTGA AAGACACCAT CAAGGTCGGC GCTGGCGAAG TCTTTGACGG 1140  
 CCACGGCGCA ACCTTCACTG CCGACAAATC TATGGGTAAC GGAGACCAGG GCGAAAATCA 1200  
 GAAGCCCATG TTCGAGCTGG CTGAAGGCGC TACGTTGAAG AATGTGAACC TGGGTGAGAA 1260  
 CGAGGTCGAT GGCATCCACG TGAAAGCCAA AAACGCTCAG GAAGTCACCA TTGACAACGT 1320  
 GCATGCCCAG AACGTCGGTG AAGACCTGAT TACGGTCAAA GGCAGGGGAG GCGCAGCGGT 1380  
 CACTAATCTG AACATCAAGA ACAGCAGTGC CAAAGGTGCA GACGACAAGG TTGTCCAGCT 1440  
 CAACGCCAAC ACTCACTTGA AAATCGACAA CTTCAAGGCC GACGATTTTCG GCACGATGGT 1500  
 TCGCACCAAC GGTGGCAAGC AGTTTGATGA CATGAGCATC GAGCTGAACG GCATCGAAGC 1560  
 TAACCACGGC AAGTTTCGCC TGGTGAAAAG CGACAGTGAC GATCTGAAGC TGGCAACGGG 1620  
 CAACATCGCC ATGACCGACG TCAAACACGC CTACGATAAA ACCCAGGCAT CGACCCAACA 1680  
 CACCGAGCTT TGAATCCAGA CAAGTAGCTT GAAAAAAGGG GGTGGACTC 1729

This DNA molecule is known as the dspE gene for *Pseudomonas syringae*. This isolated DNA molecule of the present invention encodes a protein or polypeptide which elicits a plant pathogen's hypersensitive response having an amino acid sequence of SEQ ID NO: 34 as follows:

Met	Ser	Ile	Gly	Ile	Thr	Pro	Arg	Pro	Gln	Gln	Thr	Thr	Thr	Pro	Leu
1				5					10					15	
Asp	Phe	Ser	Ala	Leu	Ser	Gly	Lys	Ser	Pro	Gln	Pro	Asn	Thr	Phe	Gly
			20					25					30		
Glu	Gln	Asn	Thr	Gln	Gln	Ala	Ile	Asp	Pro	Ser	Ala	Leu	Leu	Phe	Gly
		35					40					45			

Ser Asp Thr Gln Lys Asp Val Asn Phe Gly Thr Pro Asp Ser Thr Val  
 50 55 60  
 Gln Asn Pro Gln Asp Ala Ser Lys Pro Asn Asp Ser Gln Ser Asn Ile  
 65 70 75 80  
 Ala Lys Leu Ile Ser Ala Leu Ile Met Ser Leu Leu Gln Met Leu Thr  
 85 90 95  
 Asn Ser Asn Lys Lys Gln Asp Thr Asn Gln Glu Gln Pro Asp Ser Gln  
 100 105 110  
 Ala Pro Phe Gln Asn Asn Gly Gly Leu Gly Thr Pro Ser Ala Asp Ser  
 115 120 125  
 Gly Gly Gly Gly Thr Pro Asp Ala Thr Gly Gly Gly Gly Asp Thr  
 130 135 140  
 Pro Ser Ala Thr Gly Gly Gly Gly Gly Asp Thr Pro Thr Ala Thr Gly  
 145 150 155 160  
 Gly Gly Gly Ser Gly Gly Gly Gly Thr Pro Thr Ala Thr Gly Gly Gly  
 165 170 175  
 Ser Gly Gly Thr Pro Thr Ala Thr Gly Gly Gly Glu Gly Gly Val Thr  
 180 185 190  
 Pro Gln Ile Thr Pro Gln Leu Ala Asn Pro Asn Arg Thr Ser Gly Thr  
 195 200 205  
 Gly Ser Val Ser Asp Thr Ala Gly Ser Thr Glu Gln Ala Gly Lys Ile  
 210 215 220  
 Asn Val Val Lys Asp Thr Ile Lys Val Gly Ala Gly Glu Val Phe Asp  
 225 230 235 240  
 Gly His Gly Ala Thr Phe Thr Ala Asp Lys Ser Met Gly Asn Gly Asp  
 245 250 255  
 Gln Gly Glu Asn Gln Lys Pro Met Phe Glu Leu Ala Glu Gly Ala Thr  
 260 265 270  
 Leu Lys Asn Val Asn Leu Gly Glu Asn Glu Val Asp Gly Ile His Val  
 275 280 285  
 Lys Ala Lys Asn Ala Gln Glu Val Thr Ile Asp Asn Val His Ala Gln  
 290 295 300  
 Asn Val Gly Glu Asp Leu Ile Thr Val Lys Gly Glu Gly Gly Ala Ala  
 305 310 315 320  
 Val Thr Asn Leu Asn Ile Lys Asn Ser Ser Ala Lys Gly Ala Asp Asp  
 325 330 335  
 Lys Val Val Gln Leu Asn Ala Asn Thr His Leu Lys Ile Asp Asn Phe  
 340 345 350  
 Lys Ala Asp Asp Phe Gly Thr Met Val Arg Thr Asn Gly Gly Lys Gln  
 355 360 365

AS  
Cont'd

Phe	Asp	Asp	Met	Ser	Ile	Glu	Leu	Asn	Gly	Ile	Glu	Ala	Asn	His	Gly
370						375					380				
Lys	Phe	Ala	Leu	Val	Lys	Ser	Asp	Ser	Asp	Asp	Leu	Lys	Leu	Ala	Thr
385					390				395						400
Gly	Asn	Ile	Ala	Met	Thr	Asp	Val	Lys	His	Ala	Tyr	Asp	Lys	Thr	Gln
				405					410					415	
Ala	Ser	Thr	Gln	His	Thr	Glu	Leu								
			420												

This protein or polypeptide is about 42.9 kDa.

Please replace the paragraph beginning on page 28, line 8, and ending on page 30, line 11, with the following replacement paragraph:

The hypersensitive response elicitor polypeptide or protein derived from *Pseudomonas solanacearum* has an amino acid sequence corresponding to SEQ ID NO: 35 as follows:

AA

Met	Ser	Val	Gly	Asn	Ile	Gln	Ser	Pro	Ser	Asn	Leu	Pro	Gly	Leu	Gln
1				5					10					15	
Asn	Leu	Asn	Leu	Asn	Thr	Asn	Thr	Asn	Ser	Gln	Gln	Ser	Gly	Gln	Ser
			20					25					30		
Val	Gln	Asp	Leu	Ile	Lys	Gln	Val	Glu	Lys	Asp	Ile	Leu	Asn	Ile	Ile
		35					40					45			
Ala	Ala	Leu	Val	Gln	Lys	Ala	Ala	Gln	Ser	Ala	Gly	Gly	Asn	Thr	Gly
		50				55					60				
Asn	Thr	Gly	Asn	Ala	Pro	Ala	Lys	Asp	Gly	Asn	Ala	Asn	Ala	Gly	Ala
65					70				75					80	
Asn	Asp	Pro	Ser	Lys	Asn	Asp	Pro	Ser	Lys	Ser	Gln	Ala	Pro	Gln	Ser
				85					90					95	
Ala	Asn	Lys	Thr	Gly	Asn	Val	Asp	Asp	Ala	Asn	Asn	Gln	Asp	Pro	Met
			100					105					110		
Gln	Ala	Leu	Met	Gln	Leu	Leu	Glu	Asp	Leu	Val	Lys	Leu	Leu	Lys	Ala
		115					120					125			
Ala	Leu	His	Met	Gln	Gln	Pro	Gly	Gly	Asn	Asp	Lys	Gly	Asn	Gly	Val
		130				135					140				
Gly	Gly	Ala	Asn	Gly	Ala	Lys	Gly	Ala	Gly	Gly	Gln	Gly	Gly	Leu	Ala
145					150					155				160	

Glu Ala Leu Gln Glu Ile Glu Gln Ile Leu Ala Gln Leu Gly Gly Gly  
 165 170 175  
 Gly Ala Gly Ala Gly Gly Ala Gly Gly Gly Val Gly Gly Ala Gly Gly  
 180 185 190  
 Ala Asp Gly Gly Ser Gly Ala Gly Gly Ala Gly Gly Ala Asn Gly Ala  
 195 200 205  
 Asp Gly Gly Asn Gly Val Asn Gly Asn Gln Ala Asn Gly Pro Gln Asn  
 210 215 220  
 Ala Gly Asp Val Asn Gly Ala Asn Gly Ala Asp Asp Gly Ser Glu Asp  
 225 230 235 240  
 Gln Gly Gly Leu Thr Gly Val Leu Gln Lys Leu Met Lys Ile Leu Asn  
 245 250 255  
 Ala Leu Val Gln Met Met Gln Gln Gly Gly Leu Gly Gly Gly Asn Gln  
 260 265 270  
 Ala Gln Gly Gly Ser Lys Gly Ala Gly Asn Ala Ser Pro Ala Ser Gly  
 275 280 285  
 Ala Asn Pro Gly Ala Asn Gln Pro Gly Ser Ala Asp Asp Gln Ser Ser  
 290 295 300  
 Gly Gln Asn Asn Leu Gln Ser Gln Ile Met Asp Val Val Lys Glu Val  
 305 310 315 320  
 Val Gln Ile Leu Gln Gln Met Leu Ala Ala Gln Asn Gly Gly Ser Gln  
 325 330 335  
 Gln Ser Thr Ser Thr Gln Pro Met  
 340

It is encoded by a DNA molecule having a nucleotide sequence corresponding to SEQ ID NO: 36 as follows:

ATGTCAGTCG GAAACATCCA GAGCCCGTCG AACCTCCCGG GTCTGCAGAA CCTGAACCTC 60  
 AACACCAACA CCAACAGCCA GCAATCGGGC CAGTCCGTGC AAGACCTGAT CAAGCAGGTC 120  
 GAGAAGGACA TCCTCAACAT CATCGCAGCC CTCGTGCAGA AGGCCGCACA GTCGGCGGGC 180  
 GGCAACACCG GTAACACCGG CAACGCGCCG GCGAAGGACG GCAATGCCAA CGCGGGCGCC 240  
 AACGACCCGA GCAAGAACGA CCCGAGCAAG AGCCAGGCTC CGCAGTCGGC CAACAAGACC 300  
 GGCAACGTCG ACGACGCCAA CAACCAGGAT CCGATGCAAG CGCTGATGCA GCTGCTGGAA 360  
 GACCTGGTGA AGCTGCTGAA GGCGGCCCTG CACATGCAGC AGCCCGGCGG CAATGACAAG 420  
 GGCAACGGCG TGGGCGGTGC CAACGGCGCC AAGGGTGCCG GCGGCCAGGG CGGCCTGGCC 480  
 GAAGCGCTGC AGGAGATCGA GCAGATCCTC GCCCAGCTCG GCGGCGGCGG TGCTGGCGCC 540  
 GGCGGCGCGG GTGGCGGTGT CGGCGGTGCT GGTGGCGCGG ATGGCGGCTC CGGTGCGGGT 600



GGCGCAGGCG GTGCGAACGG CGCCGACGGC GGCAATGGCG TGAACGGCAA CCAGGCGAAC 660  
 GGCCCGCAGA ACGCAGGCGA TGTCAACGGT GCCAACGGCG CGGATGACGG CAGCGAAGAC 720  
 CAGGGCGGCC TCACCGGCGT GCTGCAAAAG CTGATGAAGA TCCTGAACGC GCTGGTGCAG 780  
 ATGATGCAGC AAGGCGGCCT CGGCGGCGGC AACCAGGCGC AGGGCGGCTC GAAGGGTGCC 840  
 GGCAACGCCT CGCCGGCTTC CGGCGCGAAC CCGGGCGCGA ACCAGCCCGG TTCGGCGGAT 900  
 GATCAATCGT CCGGCCAGAA CAATCTGCAA TCCCAGATCA TGGATGTGGT GAAGGAGGTC 960  
 GTCCAGATCC TGCAGCAGAT GCTGGCGGCG CAGAACGGCG GCAGCCAGCA GTCCACCTCG 1020  
 ACGCAGCCGA TGTA 1035

Further information regarding the hypersensitive response elicitor polypeptide or protein derived from *Pseudomonas solanacearum* is set forth in Arlat, M., F. Van Gijsegem, J. C. Huet, J. C. Pemollet, and C. A. Boucher, "PopA1, a Protein which Induces a Hypersensitive-like Response in Specific Petunia Genotypes, is Secreted via the Hrp Pathway of *Pseudomonas solanacearum*," EMBO J. 13:543-533 (1994), which is hereby incorporated by reference.

Please replace the paragraph beginning on page 30, line 12, and ending on page 30, line 25, with the following replacement paragraph:

The hypersensitive response elicitor polypeptide or protein from *Xanthomonas campestris* pv. *glycines* has an amino acid sequence corresponding to SEQ ID NO: 37 as follows:

Thr Leu Ile Glu Leu Met Ile Val Val Ala Ile Ile Ala Ile Leu Ala  
 1 5 10 15  
 Ala Ile Ala Leu Pro Ala Tyr Gln Asp Tyr  
 20 25

This sequence is an amino terminal sequence having only 26 residues from the hypersensitive response elicitor polypeptide or protein of *Xanthomonas campestris* pv. *glycines*. It matches with fimbrial subunit proteins determined in other *Xanthomonas campestris* pathovars.

Please replace the paragraph beginning on page 30, line 26, and ending on page 30, line 34, with the following replacement paragraph:

AC<sup>0</sup>  
The hypersensitive response elicitor polypeptide or protein from *Xanthomonas campestris* pv. *pelargonii* is heat stable, protease sensitive, and has a molecular weight of 20 kDa. It includes an amino acid sequence corresponding to SEQ ID NO: 38 as follows:

Ser	Ser	Gln	Gln	Ser	Pro	Ser	Ala	Gly	Ser	Glu	Gln	Gln	Leu	Asp	Gln
1				5					10					15	
<hr/>															
Leu	Leu	Ala	Met												
			20												

Please replace the paragraph beginning on page 32, line 27, and ending on page 33, line 5, with the following replacement paragraph:

g/b  
A11  
~~An example of suitable fragments of a hypersensitive response elicitor which~~  
do not elicit a hypersensitive response include fragments of the *Erwinia amylovora* hypersensitive response elicitor. Suitable fragments include a C-terminal fragment of the amino acid sequence of SEQ ID NO: 23, an N-terminal fragment of the amino acid sequence of SEQ ID NO: 23, or an internal fragment of the amino acid sequence of SEQ ID NO: 23. The C-terminal fragment of the amino acid sequence of SEQ ID NO: 23 can span the following amino-acids of SEQ ID NO: 23: 169 and 403, 210 and 403, 267 and 403, or 343 and 403. The internal fragment of the amino acid sequence of SEQ ID NO: 23 can span the following amino acids of SEQ ID NO: 23: 105 and 179, 137 and 166, 121 and 150, or 137 and 156. ~~Other suitable fragments can be identified in accordance with the present invention.~~

Please replace the paragraph beginning on page 33, line 6, and ending on page 33, line 10, with the following replacement paragraph:

A12  
Another example of a useful fragment of a hypersensitive response elicitor which fragment does not itself elicit a hypersensitive response is the protein fragment containing amino acids 190 to 294 of the amino acid sequence (SEQ ID NO: 31) for the *Pseudomonas syringae* pv. *syringae* hypersensitive response elicitor. This fragment is useful in imparting disease resistance and enhancing plant growth.

Please replace the paragraph beginning on page 33, line 11, and ending on page 33, line 14, with the following replacement paragraph:

AB  
Yet another example of a useful fragment of a hypersensitive response elicitor is the peptide having an amino acid sequence corresponding to SEQ ID NO: 39. This peptide is derived from the hypersensitive response eliciting glycoprotein of *Phytophthora megasperma* and enhances plant growth.

Please replace the paragraph beginning on page 46, line 29, and ending on page 47, line 12, with the following replacement paragraph:

ALY  
*Escherichia coli* BL21(DE3) strains containing the hrpN clones were grown in Luria broth medium (5g/L Difco Yeast extract, 10 g/L Difco Tryptone, 5 g/L NaCl, and 1 mM NaOH) containing 30 µg/ml of kanamycin at 37°C overnight. The bacteria were then inoculated into 100 volumes of the same medium and grown at 37°C to an OD<sub>620</sub> of 0.6-0.8. The bacteria were then inoculated into 250 volumes of the same medium and grown at 37°C to an OD<sub>620</sub> of ca. 0.3 or 0.6-0.8. One milli molar IPTG was then added and the cultures grown at 19°C overnight (ca. 18 hours). Not all of the clones were successfully expressed using this strategy. Several of the clones had to be grown in Terrific broth (12 g/L Bacto Tryptone, 24 g/L Bacto yeast, 0.4% glycerol, 0.17 M KH<sub>2</sub>PO<sub>4</sub>, and 0.72 K<sub>2</sub>HPO<sub>4</sub>), and/or grown at 37°C after IPTG induction, and/or harvested earlier than overnight (Table 1).

Table 1: Expression of hypersensitive response elicitor truncated proteins

Fragment	amino acids (SEQ ID NO: 23)	Growth medium	Induction O.D.	Expression temp.	Harvest time
1 (+ control)	1-403	LB	ca. 0.3 or 0.6-0.8	19°C or 25°C	16-18 hr
2 (+ control)	-	LB and TB	ca. 0.3 or 0.6-0.8	19 C and 37 C	16-18 hr
3	105-403	LB	0.6-0.8	19°C	16-18 hr
4	169-403	TB	ca. 0.3	19°C	16-18 hr
5	210-403	LB or M9ZB	0.6-0.8	19°C	16-18 hr
6	257-403	LB or M9ZB	0.6-0.8	19°C	16-18 hr
7	343-403	LB	ca. 0.3	19°C	5 hr
8	1-75	TB	ca. 0.3	37°C	16-18 hr
9	1-104	TB	ca. 0.3	37°C	16-18 hr
10	1-168	TB	ca. 0.3	37°C	16-18 hr
11	1-266	LB	ca. 0.3	37°C	4 hr
12	1-342	LB	0.6-0.8	19°C	16-18 hr
13	76-209	LB	ca. 0.3	37°C	5 hr
14	76-168	TB or LB	ca. 0.3	37°C	3 hr or 16-18 hr
15	105-209	M9ZB	ca. 0.3	37°C	3 hr
16	169-209	no expression			
17	105-168	LB	ca. 0.3	37°C	3-5 hr
18	99-209	LB	ca. 0.3	37°C	3 hr
19	137-204	LB	ca. 0.3	37°C	3 hr
20	137-180	LB	ca. 0.3	37°C	16-18 hr.
21	105-180	LB	ca. 0.3	37°C	3 hr
22	150-209	no expression			
23	150-180	no expression			

Please replace the paragraph beginning on page 51, line 2, and ending on page 52, line 2, with the following replacement paragraph:

715 The small scale expression and purification of the fragment proteins was done to screen for expression and HR activity (Table 2).

Table 2

Expression and HR activity of hypersensitive response elicitor truncated proteins (small scale screening)

Fragment #	Amino Acids (SEQ ID NO: 23)	Expression	HR activity
1(+control)	1-403	+	+
2(- control)	-	background protein only	-
3	105-403	+	+
4	169-403	+	-
5	210-403	+	-
6	267-403	+	-
7	343-403	+/-	-
8	1-75	+	-
9	1-104	+	+/-
10	1-168	+	+
11	1-266	+	+
12	1-342	+	+
13	76-209	+	+
14	76-168	+	-
15	105-209	+	+
16	169-209	-	-
17	105-168	+	-
18	99-209	+	+
19	137-204	+	+
20	137-180	+	+
21	105-180	+	+
22	150-209	-	-
23	150-180	-	-

All of the cloned fragment proteins were expressed at varying levels except for three small fragments (amino acids 169-209, 150-209, and 150-180). Fragments 210-403 and 267-403 were expressed very well, yielding a high concentration of protein from a small scale purification, resulting in a substantial protein band on SDS gel electrophoresis. Other fragments (such as a.a. 1-168 and 1-104) produced much less protein, resulting in faint protein bands upon electrophoresis. It was difficult to determine whether fragment 343-403, the smallest C-terminal protein, was expressed, as there were several background proteins apparent on the gel, in addition to the suspected 343-403 protein. The positive and negative

AB control proteins, consisting of the full length hypersensitive response elicitor protein and only background proteins, respectively, were tested for expression and HR activity as well.

Please replace the paragraph beginning on page 52, line 3, and ending on page 53, line 2, with the following replacement paragraph:

The large scale expression and purification of the fragment proteins was done to determine the level of expression and titer of the HR activity (Table 3).

Table 3

Expression level and HR titer of hypersensitive response elicitor truncated proteins (large scale purification)

Fragment #	Amino acids (SEQ ID NO: 23)	Expression	HR titer
1(+ control)	1-403	3.7 mg/ml	5-7 µg/ml
2(- control)	-	-	1:2 dilution
4	169-403	2 mg/ml	-
5	210-403	5 mg/ml	-
6	267-403	4 mg/ml	-
7	343-402	200µg/ml	-
8	1-75	50µg/ml	-
9	1-104	50µg/ml	3 µg/ml (1:16 dilution)
10	1-168	1 mg/ml	1 µg/ml
13	76-209	2.5 mg/ml	5 µg/ml
14	76-168	2 mg/ml	-
15	105-209	5 mg/ml	5-10µg/ml
17	105-168	250µg/ml	-
19	137-204	3.6 mg/ml	3.5 µg/ml
20	137-180	250 µg/ml	16 µg/ml

The truncated proteins deemed to be the most important in characterizing the hypersensitive response elicitor were chosen for large scale expression. The positive control (full length hypersensitive response elicitor) was expressed at a relatively high level at 3.7 mg/ml. All of the C-terminal proteins were expressed at relatively high levels from 2-5 mg/ml, except for fragment 343-403 as discussed earlier. The N-terminal fragments were expressed very well also; however, during the purification process, the protein precipitated and very little was resolubilized. The concentrations in Table 3 reflect only the solubilized protein. The internal fragments were expressed in the range of 2-3.6 mg/ml. It was extremely difficult to determine the concentration of fragment 105-168 (it was suspected that the concentration was

446 much higher than indicated), as the protein bands on the SDS gel were large, but poorly stained. The negative control contained several background proteins as expected, but no obviously induced dominant protein.

Please replace the paragraph beginning on page 53, line 14, and ending on page 53, line 26, with the following replacement paragraph:

A17 One definitive domain encompassing a small internal region of the protein from a.a. 137-180 (SEQ ID NO: 23), a mere 44 a.a, is identified as the smallest HR domain. The other potential HR domain is thought to be located in the N-terminus of the protein from a.a. 1-104 (possibly a.a. 1-75) (SEQ ID NO: 23). It was difficult to confirm or narrow down the N-terminus HR domain due to the difficulties encountered in purifying these fragment proteins. The N-terminus fragment proteins had to be purified with urea as no protein was recovered when the native purification process was used. Consequently, these proteins precipitated during the renaturing process and were difficult or nearly impossible to get back into solution, thereby making it hard to run the proteins through the HR assay, as only soluble protein is able to elicit HR. Difficulty narrowing the N-terminus HR domain was only compounded by the fact that the negative control elicited false HR at the low dilution levels thereby reducing the sensitivity of the assay.

Please replace the paragraph beginning on page 53, line 27, and ending on page 53, line 33, with the following replacement paragraph:

A18 Surprisingly, when the internal HR domain was cleaved between a.a. 168 and 169 (fragments 76-168 and 105-168) (SEQ ID NO: 23) the fragment lost its HR activity. This suggests that the HR activity of fragment 1-168 (SEQ ID NO: 23) should not be attributed to the internal HR domain, but rather to some other domain, leading to the assumption that there was likely a second HR domain to be found in the N-terminal region of the protein. However, as discussed earlier it was difficult to confirm this assumption.

Please replace the paragraph beginning on page 54, line 1, and ending on page 54, line 6, with the following replacement paragraph:

AM The hypersensitive response elicitor C-terminus (a.a. 210-403 (SEQ ID NO: 23)) did not contain an HR domain. It did not elicit HR at a detectable level using the current HR assay. Even the large C-terminal fragment from a.a. 169-403 (SEQ ID NO: 23) did not elicit HR even though it contained part of the internal HR domain. As stated above, cleaving the protein between amino acids 168 and 169 (SEQ ID NO: 23) causes a loss of HR activity.

Please replace the paragraph beginning on page 54, line 7, and ending on page 54, line 18, with the following replacement paragraph:

A<sup>20</sup> Because some of the small cloned proteins with 61 a.a. or less were not expressed, several oligopeptides were synthesized with 30 a.a. to narrow down the functional region of the internal HR domain. The oligopeptides were synthesized within the range of a.a. 121-179 (SEQ ID NO: 23). However, these oligos did not elicit HR. It was not expected that there would be an HR from oligos 137-166, 121-150, and 137-156 (SEQ ID NO: 23) as these fragments did not contain the imperative amino acids 168 and 169 (SEQ ID NO: 23). It was expected that the oligo 150-179 (SEQ ID NO: 23) would elicit an HR. It is possible that 30 a.a. is too small for the protein to elicit any activity due to a lack of folding and, therefore, a lack of binding or that during the synthesis of the peptides important amino acids were missed (either in the process, or simply by the choice of which 30 amino acids to synthesize) and, therefore, the fragments would not be able to elicit HR.

Please replace the paragraph beginning on page 59, line 19, and ending on page 60, line 2, with the following replacement paragraph:

A<sup>21</sup> To test if the 13 amino acid peptide derived from the 42 kDa protein also enhanced plant growth, 20 mg of the oligopeptide was synthesized from Biosynthesis Corp. The synthesized sequence of the peptide is NH<sub>2</sub>-Val-Trp-Asn-Gln-Pro-Val-Arg-Gly-Phe-Lys-Val-Tyr-Glu-COOH (SEQ ID NO: 39). The synthesized peptide was resuspended in 10 ml of 5 mM potassium phosphate buffer and, then, diluted to 1 and 100 ng/ml with the same